ARDAC Kidney Health Study



Screening Report

Mt Druitt

2010





The ARDAC Study

The ARDAC Study (Antecedents of Renal Disease in Aboriginal Children) is based at The Centre for Kidney Research at The Children's Hospital at Westmead and is investigating the early signs of kidney and cardiovascular disease in both Aboriginal and non-Aboriginal children in NSW. The first phase of the study was conducted over 6 years between 2002 and 2007 and followed the health of more than 2000 primary school children from areas across NSW. Initially participants were recruited from primary schools, however they are now screened in high schools, at home and in the community as the participants are older and some have left school. Screening in Phase 2 of the study includes measurement of height, weight, blood pressure, waist circumference and a urine test.

The results of the first phase showed no difference in presence of markers for kidney and cardiovascular disease between Aboriginal and non-Aboriginal participants. Screening for Phase 2 started in 2008. The success of the ARDAC study has been the result of the support and commitment from Aboriginal Health Workers (AHW's), Aboriginal Education Officers (AEO's), teachers, staff from the Aboriginal Medical Services, the Division of General Practice and a large number of participating schools and communities.


In 2010 we screened 68 participants in the Mt Druitt Area. These are the results of screening in the Mt Druitt Area.


Schools screened in Mt Druitt Area:

School	Participants
Blacktown Girls HS	2
Cambridge Park HS	1
Chifley College – Bidwill Campus	12
Chifley College – Dunheved Campus	10
Chifley College – Mt Druitt Campus	4
Chifley College – Senior Campus	3
Chifley College – Shalvey Campus	3
Clare Catholic HS	1
Kellyville HS	2
Mitchell HS	1
Penrith HS	4
Plumpton HS	3
St Marys Senior College	2
The Hills Sports HS	2
Home Visits	18
Total	68

ARDAC Phase 2 Screening Report

Mt Druitt Area 2010





The first graph shows the number of Aboriginal and non-Aboriginal participants who were referred for each risk factor. The second graph shows the percentage of Aboriginal and non-Aboriginal participants who were referred for each risk factor.

Number of participants tested: 68

Ethnicity: 50% non-Aboriginal (34) and 50% Aboriginal (34)

■ Blood in urine \geq 1+ (25 cells/uL) excluding menstruation:

There was 0 out of 68 participants (0%) who had blood in their urine, 0 non-Aboriginal and 0 Aboriginal

Protein in urine ≥ 1+ (30mg/dL):

There were 20 out of 68 participants (29%) with protein in their urine, 13 non-Aboriginal and 7 Aboriginal

■ High blood pressure (systolic blood pressure > 95% for age):

There were 6 out of 68 participants (9%) with systolic hypertension, 4 non-Aboriginal and 2 Aboriginal

■ Obesity (Body Mass Index SDS ≥2):

There were 7 out of 68 participants (10%) with a BMI SDS \geq 2, 4 non-Aboriginal and 3 Aboriginal

Referral

Participants are referred to their local doctor, Aboriginal Medical Service or Community Health Centre for further investigation if one or more of the following are found during screening;

- High blood pressure
- Obesity
- Blood, protein or albumin in their urine

Participants who require follow-up receive instructions with their screening results.

Contact Us

If you have any questions about screening or results please contact us on:

Free call: 1800 005 846 Phone: 9845 1469 Fax: 9845 1491

Email: nicolav@chw.edu.au or rachaek2@chw.edu.au