## ARDAC Kidney Health Study



# Screening Report Far West NSW 2014





### The ARDAC Study

The ARDAC Study (Antecedents of Renal Disease in Aboriginal Children) is based at The Centre for Kidney Research at The Children's Hospital at Westmead and is investigating the early signs of kidney and cardiovascular disease in both Aboriginal and non-Aboriginal children in NSW. The first phase of the study was conducted over 6 years between 2002 and 2007 and followed the health of more than 2000 primary school children from areas across NSW. Initially participants were recruited from primary schools, however they are now screened in high schools, at home and in the community as the participants are older and some have left school. Screening in Phase 2 of the study includes measurement of height, weight, blood pressure, waist circumference and a urine test.

The results of the first phase showed no difference in presence of markers for kidney and cardiovascular disease between Aboriginal and non-Aboriginal participants. Screening for Phase 2 started in 2008. The success of the ARDAC study has been the result of the support and commitment from Aboriginal Health Workers (AHW's), Aboriginal Education Officers (AEO's), teachers, staff from the Aboriginal Medical Services, the Division of General Practice and a large number of participating schools and communities.


In 2014 we screened 114 participants from Broken Hill, Wilcannia and Menindee. These are the results of screening in Broken Hill, Wilcannia and Menindee in 2014.


Screening locations:

School	Participants
Broken Hill High School	43
Menindee Central School	33
Menindee Health Service	3
Wilcannia Central School	4
Wilcannia Health Service	13
Willyama High School	2
Home Visits	16
Total	114

#### **ARDAC Phase 2 Screening Report**

Far West NSW 2014





The first graph shows the number of Aboriginal and non-Aboriginal participants who were referred for each risk factor. The second graph shows the percentage of Aboriginal and non-Aboriginal participants who were referred for each risk factor.

#### Number of participants tested: 114

Ethnicity: 39.5% non-Aboriginal (45) and 60.5% Aboriginal (69)

#### • Blood in urine $\geq$ 1+ (25 cells/uL) excluding menstruation:

There were 5 out of 114 participants (4%) who had blood in their urine, 1 non-Aboriginal and 4 Aboriginal

#### Protein in urine <a>> 1+ (30mg/dL):</a>

There were 30 out of 114 participants (26%) with protein in their urine, 9 non-Aboriginal and 21 Aboriginal

#### • High blood pressure (systolic blood pressure > 95% for age ):

There were 18 out of 114 participants (16%) with systolic hypertension, 9 non-Aboriginal and 9 Aboriginal

#### Obesity (Body Mass Index SDS <u>></u>2):

There were 27 out of 114 participants (24%) with a BMI SDS  $\geq$  2, 8 non-Aboriginal and 19 Aboriginal

#### <u>Referral</u>

Participants are referred to their local doctor, Royal Flying Doctor Service or Aboriginal Medical Service for further investigation if one or more of the following are found during screening;

- High blood pressure
- Obesity
- Blood, protein or albumin in their urine

Participants who require follow-up receive instructions with their screening results.

#### <u>Contact Us</u>

If you have any questions about screening or results please contact us on:

Free call: 1800 005 846 Phone: 9845 1469 Fax: 9845 1491

Email: nicola.vukasin@health.nsw.gov.au or rachael.kearns@health.nsw.gov.au