

ARDAC Kidney Health Study



Screening Report

Goulburn & Queanbeyan

23rd – 27th March 2009

ckr

CENTRE FOR
KIDNEY RESEARCH
The Children's Hospital
at Westmead



The ARDAC Study

The ARDAC Study (**A**ntecedents of **R**enal **D**isease in **A**boriginal **C**hildren) is based at The Centre for Kidney Research at The Children's Hospital at Westmead and is investigating the early signs of kidney and cardiovascular disease in both Aboriginal and non-Aboriginal children in NSW. The first phase of the study was conducted over 6 years between 2002 and 2007 and followed the health of more than 2000 primary school children from areas across NSW. Initially participants were recruited from primary schools, however they are now screened in high schools, at home and in the community as the participants are older and some have left school. Screening in Phase 2 of the study includes measurement of height, weight, blood pressure, waist circumference and a urine test.



The results of the first phase showed no difference in presence of markers for kidney and cardiovascular disease between Aboriginal and non-Aboriginal participants. Screening for Phase 2 started in 2008. The success of the ARDAC study has been the result of the support and commitment from Aboriginal Health Workers (AHW's), Aboriginal Education Officers (AEO's), teachers, staff from the Aboriginal Medical Services, the Division of General Practice and a large number of participating schools and communities.

In 2009 we screened 69 participants from Goulburn and Queanbeyan. These are the results of screening in Goulburn and Queanbeyan in 2009.

Schools screened in the Goulburn and Queanbeyan:

School	Participants
Bradfordville Public School	1
Goulburn High School	11
Goulburn West Public School	1
Karabar High School	20
Mulwaree High School	8
Queanbeyan High School	14
Queanbeyan South Public School	3
Queanbeyan West Public School	1
Trinity Catholic College	1
Home Visits	9
Total	69

ARDAC Phase 2 Screening Report Goulburn & Queanbeyan



The first graph shows the number of Aboriginal and non-Aboriginal participants who were referred for each risk factor. The second graph shows the percentage of Aboriginal and non-Aboriginal participants who were referred for each risk factor.

Number of participants tested: 69

Ethnicity: 41% non-Aboriginal (28) and 59% Aboriginal (41)

▪ **Blood in urine \geq 1+ (25 cells/uL) excluding menstruation:**

There were 2 out of 69 participants (3%) who had blood in their urine, 1 non-Aboriginal and 1 Aboriginal

▪ **Protein in urine \geq 1+ (30mg/dL):**

There were 12 out of 69 participants (17%) with protein in their urine, 4 non-Aboriginal and 8 Aboriginal

▪ **High blood pressure (systolic blood pressure > 95% for age):**

There were 2 out of 69 participants (3%) with systolic hypertension, 0 non-Aboriginal and 2 Aboriginal

▪ **Obesity (Body Mass Index SDS \geq 2):**

There were 5 out of 69 participants (7%) with a BMI SDS \geq 2, 1 non-Aboriginal and 4 Aboriginal

Referral

Participants are referred to their local doctor, Aboriginal Medical Service or Community Health Centre for further investigation if one or more of the following are found during screening;

- **High blood pressure**
- **Obesity**
- **Blood, protein or albumin in their urine**

Participants who require follow-up receive instructions with their screening results.

Contact Us

If you have any questions about screening or results please contact us on:

Free call: 1800 005 846

Phone: 9845 1469

Fax: 9845 1491

Email: ardac@chw.edu.au